

SPECIAL REPORT

This Special Report is brought to you by:

keith deary

Stock Market Forex Guides Tools and Articles

For More Details, Visit Us At:

<http://www.forexnstocks.com>

Copyright 2008 <http://Traffic-Sales-Profits.com>

Legal Notice:

The Publisher has strived to be as accurate and complete as possible in the creation of this book, notwithstanding the fact that he does not warrant or represent at any time that the contents within are accurate due to the rapidly changing nature of the Internet. While all attempts have been made to verify information provided in this publication, the Publisher assumes no responsibility for errors, omissions, or contrary interpretation of the subject matter herein. Any perceived slights of specific persons, peoples, or organizations are unintentional. In practical manuals, like anything else in life, there are no guarantees of income made. Readers are cautioned to rely on their own judgment about their individual circumstances to act accordingly. This book is not intended for use as a source of legal, business, accounting or financial advice. All readers are advised to seek services of competent professionals in legal, business, accounting, and finance areas.

SPECIAL REPORT:

GARDENING

This special report is brought to you by:

<http://Traffic-Sales-profits.com/>

**I Just Figured Out How To Attract
FREE Traffic To My Website... Have
You?**

**Did You Know There Are 15 Ways To
Become A "TRAFFIC MAGNET"?**

**Click Here Now To Read How I Got
67115 Visitors This Month...**

Visit <http://Traffic-Sales-Profits.com>



Limits of Liability/Disclaimer of Warranty

The author and publisher of this book and the accompanying materials have used their best efforts in preparing this report. The author and publisher make no representation or warranties with respect to the accuracy, applicability, fitness, or completeness of the contents of this program.

They disclaim any warranties (expressed or implied), merchantability, or fitness for any particular purpose. The author and publisher shall no event be held liable for any loss or other damages, including but not limited to special, incidental, consequential or other damages. As always, the advice of a competent legal, tax, accounting or other professionals should be sought.

All information is for educational purpose only and is not warranted for content, accuracy or any other implied or explicit purpose. The information presented represents the views of the author as of the date of publication. Because of the rate with which conditions change, the author reserve the rights to alter and update their opinions based on the new conditions. This report is for informational purposes only and the author does not accept any responsibilities for any liabilities resulting from the use of this information. While every attempt has been made to verify the information provided here, the author and his resellers and affiliates cannot assume any responsibility for errors, inaccuracies or omissions. Any sleights of people or organizations are unintentional.

This book contains material protected under International and Federal Copyright Laws and Treaties. Any unauthorized reprint or use of this material is prohibited.

Table of Contents:

The Genesis Of Soil.....	4
Making A Garden.	6
Planting Seeds.	10
Fighting Plant Enemies.	12
Garden Pests.....	15
Landscape Gardening.	18
The Cultivation Of Vegetables.	23
Vegetable Culture.	28
Wild-Flower Garden.....	33
Recommended Resources	37

THE GENESIS OF SOIL.

Soil primarily had its beginning from rock together with animal and vegetable decay, if you can imagine long stretches or periods of time when great rock masses were crumbling and breaking up. Heat, water action, and friction were largely responsible for this. By friction here is meant the rubbing and grinding of rock mass against rock mass. Think of the huge rocks, a perfect chaos of them, bumping, scraping, and settling against one another. What would be the result? Well, I am sure you all could work that out. This is what happened: bits of rock were worn off, a great deal of heat was produced, and pieces of rock were pressed together to form new rock masses, some portions becoming dissolved in water. Why, I myself, almost feel the stress and strain of it all. Can you?

Then, too, there were great changes in temperature. First everything was heated to a high temperature, and then gradually became cool. Just think of the cracking, the crumbling, and the upheavals that such changes must have caused! You know some of the effects in winter of sudden freezes and thaws. But the little examples of bursting water pipes and broken pitchers are as nothing to what was happening in the world during those days. The water and the gases in the atmosphere helped along this crumbling work.

From all this action of rubbing, which action we call mechanical, it is easy enough to understand how sand was formed. This represents one of the great divisions of soil sandy soil. The sea shores are great masses of pure sand. If soil were nothing but broken rock masses then indeed it would be very poor and unproductive. But the early forms of animal and vegetable life decaying became a part of the rock mass and a better soil resulted. So the soils we speak of as sandy soils have mixed with the sand other matter, sometimes clay, sometimes vegetable matter or humus, and often animal waste.

Clay brings us right to another class of soils clayey soils. It happens that certain portions of rock masses became dissolved when water trickled over them and heat was plenty and abundant. This dissolution took place largely because there is in the air a certain gas called carbon dioxide or carbonic acid gas. This gas attacks and changes certain substances in rocks. Sometimes you see great rocks with portions sticking up looking as if they had been eaten away. Carbonic acid did this. It changed this eaten part into something else which we call clay. A change like this is not mechanical but chemical. The difference in the two kinds of change is just this: in the one case of sand, where a mechanical change went on, you still have just what you started with, save that the size of the mass is smaller. You started with a big rock, and ended with little particles of sand. But you had no different kind of rock in the end. Mechanical action might be illustrated with a piece of lump sugar. Let the sugar represent a big mass of rock. Break up the sugar, and even the smallest bit is sugar. It is just so with the rock mass; but in the case of a chemical change you start with one thing and end with another. You started with a big mass of rock which had in it a portion that became changed by the acid acting on it. It ended in being an entirely different thing which we call clay. So in the case of chemical change a certain something is started with and in the end we have an entirely different thing. The clay soils are often called mud soils because of the amount of water used in their formation.

The third sort of soil which we farm people have to deal with is lime soil. Remember we are thinking of soils from the farm point of view. This soil of course ordinarily was formed from limestone. Just as soon as one thing is mentioned about which we know nothing, another comes up of which we are just as ignorant. And so a whole chain of questions follows. Now you are probably saying within yourselves, how was limestone first formed?

At one time ages ago the lower animal and plant forms picked from the water particles of lime. With the lime they formed skeletons or houses about

themselves as protection from larger animals. Coral is representative of this class of skeleton-forming animal.

As the animal died the skeleton remained. Great masses of this living matter pressed all together, after ages, formed limestone. Some lime stones are still in such shape that the shelly formation is still visible. Marble, another limestone, is somewhat crystalline in character. Another well-known limestone is chalk. Perhaps you'd like to know a way of always being able to tell limestone. Drop a little of this acid on some lime. See how it bubbles and fizzes. Then drop some on this chalk and on the marble, too. The same bubbling takes place. So lime must be in these three structures. One does not have to buy a special acid for this work, for even the household acids like vinegar will cause the same result.

Then these are the three types of soil with which the farmer has to deal, and which we wish to understand. For one may learn to know his garden soil by studying it, just as one learns a lesson by study.

MAKING A GARDEN.

The first thing in garden making is the selection of a spot. Without a choice, it means simply doing the best one can with conditions. With space limited it resolves itself into no garden, or a box garden. Surely a box garden is better than nothing at all.

But we will now suppose that it is possible to really choose just the right site for the garden. What shall be chosen? The greatest determining factor is the sun. No one would have a north corner, unless it was absolutely forced upon him; because, while north corners do for ferns, certain wild flowers, and begonias, they are of little use as spots for a general garden.

If possible, choose the ideal spot a southern exposure. Here the sun lies warm all day long. When the garden is thus located the rows of vegetables and flowers should run north and south. Thus placed, the plants receive the sun's rays all the morning on the eastern side, and all the afternoon on the western side. One ought not to have any lopsided plants with such an arrangement.

Suppose the garden faces southeast. In this case the western sun is out of the problem. In order to get the best distribution of sunlight run the rows northwest and southeast.

The idea is to get the most sunlight as evenly distributed as possible for the longest period of time. From the lopsided growth of window plants it is easy enough to see the effect on plants of poorly distributed light. So if you use a little diagram remembering that you wish the sun to shine part of the day on one side of the plants and part on the other, you can juggle out any situation. The southern exposure gives the ideal case because the sun gives half time nearly to each side. A northern exposure may mean an almost entire cut-off from sunlight; while northeastern and southwestern places always get uneven distribution of sun's rays, no matter how carefully this is planned.

The garden, if possible, should be planned out on paper. The plan is a great help when the real planting time comes. It saves time and unnecessary buying of seed.

New garden spots are likely to be found in two conditions: they are covered either with turf or with rubbish. In large garden areas the ground is ploughed and the sod turned under; but in small gardens remove the sod. How to take off the sod in the best manner is the next question. Stake and line off the garden spot. The line gives an accurate and straight course to follow. Cut the edges with the spade all along the line. If the area is a small one, say four feet by eighteen or twenty,

this is an easy matter. Such a narrow strip may be marked off like a checkerboard, the sod cut through with the spade, and easily removed. This could be done in two long strips cut lengthwise of the strip. When the turf is cut through, roll it right up like a roll of carpet.

But suppose the garden plot is large. Then divide this up into strips a foot wide and take off the sod as before. What shall be done with the sod? Do not throw it away for it is full of richness, although not quite in available form. So pack the sod grass side down one square on another. Leave it to rot and to weather. When rotted it makes a fine fertilizer. Such a pile of rotting vegetable matter is called a compost pile. All through the summer add any old green vegetable matter to this. In the fall put the autumn leaves on. A fine lot of goodness is being fixed for another season.

Even when the garden is large enough to plough, I would pick out the largest pieces of sod rather than have them turned under. Go over the ploughed space, pick out the pieces of sod, shake them well and pack them up in a compost heap.

Mere spading of the ground is not sufficient. The soil is still left in lumps. Always as one spades one should break up the big lumps. But even so the ground is in no shape for planting. Ground must be very fine indeed to plant in, because seeds can get very close indeed to fine particles of soil. But the large lumps leave large spaces which no tiny root hair can penetrate. A seed is left stranded in a perfect waste when planted in chunks of soil. A baby surrounded with great pieces of beefsteak would starve. A seed among large lumps of soil is in a similar situation. The spade never can do this work of pulverizing soil. But the rake can. That's the value of the rake. It is a great lump breaker, but will not do for large lumps. If the soil still has large lumps in it take the hoe.

Many people handle the hoe awkwardly. The chief work of this implement is to rid the soil of weeds and stir up the top surface. It is used in summer to form that mulch of dust so valuable in retaining moisture in the soil. I often see people as if they were going to chop into atoms everything around. Hoeing should never be such vigorous exercise as that. Spading is vigorous, hard work, but not hoeing and raking.

After lumps are broken use the rake to make the bed fine and smooth. Now the great piece of work is done.



**Creating a Beautiful Garden Couldn't be Easier,
with this Expert Guide**

Jam-packed with 155 pages of info, The Weekend Gardener is a virtual encyclopedia of tips, tricks, techniques, and secrets of master gardeners from around

[\[Click Here To Learn More\]](#)

PLANTING SEEDS.

Any reliable seed house can be depended upon for good seeds; but even so, there is a great risk in seeds. A seed may to all appearances be all right and yet not have within it vitality enough, or power, to produce a hardy plant.

If you save seed from your own plants you are able to choose carefully. Suppose you are saving seed of aster plants. What blossoms shall you decide upon? Now it is not the blossom only which you must consider, but the entire plant. Why? Because a weak, straggly plant may produce one fine blossom. Looking at that one blossom so really beautiful you think of the numberless equally lovely plants you are going to have from the seeds. But just as likely as not the seeds will produce plants like the parent plant.

So in seed selection the entire plant is to be considered. Is it sturdy, strong, well shaped and symmetrical; does it have a goodly number of fine blossoms? These are questions to ask in seed selection.

If you should happen to have the opportunity to visit a seeds man's garden, you will see here and there a blossom with a string tied around it. These are blossoms chosen for seed. If you look at the whole plant with care you will be able to see the points which the gardener held in mind when he did his work of selection.

In seed selection size is another point to hold in mind. Now we know no way of telling anything about the plants from which this special collection of seeds came. So we must give our entire thought to the seeds themselves. It is quite evident that there is some choice; some are much larger than the others; some far plumper, too. By all means choose the largest and fullest seed. The reason is this: When you break open a bean and this is very evident, too, in the peanut you

see what appears to be a little plant. So it is. Under just the right conditions for development this 'little chap' grows into the bean plant you know so well.

This little plant must depend for its early growth on the nourishment stored up in the two halves of the bean seed. For this purpose the food is stored. Beans are not full of food and goodness for you and me to eat, but for the little baby bean plant to feed upon. And so if we choose a large seed, we have chosen a greater amount of food for the plantlet. This little plantlet feeds upon this stored food until its roots are prepared to do their work. So if the seed is small and thin, the first food supply insufficient, there is a possibility of losing the little plant.

You may care to know the name of this pantry of food. It is called a cotyledon if there is but one portion, cotyledons if two. Thus we are aided in the classification of plants. A few plants that bear cones like the pines have several cotyledons. But most plants have either one or two cotyledons.

From large seeds come the strongest plantlets. That is the reason why it is better and safer to choose the large seed. It is the same case exactly as that of weak children.

There is often another trouble in seeds that we buy. The trouble is impurity. Seeds are sometimes mixed with other seeds so like them in appearance that it is impossible to detect the fraud. Pretty poor business, is it not? The seeds may be unclean. Bits of foreign matter in with large seed are very easy to discover. One can merely pick the seed over and make it clean. By clean is meant freedom from foreign matter. But if small seed are unclean, it is very difficult, well nigh impossible, to make them clean.

The third thing to look out for in seed is viability. We know from our testing that seeds which look to the eye to be all right may not develop at all. There are reasons. Seeds may have been picked before they were ripe or mature; they

may have been frozen; and they may be too old. Seeds retain their viability or germ developing power, a given number of years and are then useless. There is a viability limit in years which differs for different seeds.

From the test of seeds we find out the germination percentage of seeds. Now if this percentage is low, don't waste time planting such seed unless it is small seed. Immediately you question that statement. Why does the size of the seed make a difference? This is the reason. When small seed is planted it is usually sown in drills. Most amateurs sprinkle the seed in very thickly. So a great quantity of seed is planted. And enough seed germinates and comes up from such close planting. So quantity makes up for quality.

But take the case of large seed, like corn for example. Corn is planted just so far apart and a few seeds in a place. With such a method of planting the matter of per cent, of germination is most important indeed.

Small seeds that germinate at fifty per cent. May be used but this is too low a per cent for the large seed. Suppose we test beans. The percentage is seventy. If low-vitality seeds were planted, we could not be absolutely certain of the seventy per cent coming up. But if the seeds are lettuce go ahead with the planting.

FIGHTING PLANT ENEMIES.

The devices and implements used for fighting plant enemies are of two sorts:

- (1) Those used to afford mechanical protection to the plants;
- (2) Those used to apply insecticides and fungicides.

Of the first the most useful is the covered frame. It consists usually of a wooden box, some eighteen inches to two feet square and about eight high, covered with glass, protecting cloth, mosquito netting or mosquito wire. The first two coverings have, of course, the additional advantage of retaining heat and protecting from cold, making it possible by their use to plant earlier than is otherwise safe. They are used extensively in getting an extra early and safe start with cucumbers, melons and the other vine vegetables.

Simpler devices for protecting newly-set plants, such as tomatoes or cabbage, from the cut-worm, are stiff, tin, cardboard or tar paper collars, which are made several inches high and large enough to be put around the stem and penetrate an inch or so into the soil.

For applying poison powders, the home gardener should supply himself with a powder gun. If one must be restricted to a single implement, however, it will be best to get one of the hand-power, compressed-air sprayers. These are used for applying wet sprays, and should be supplied with one of the several forms of mist-making nozzles, the non-cloggable automatic type being the best. For more extensive work a barrel pump, mounted on wheels, will be desirable, but one of the above will do a great deal of work in little time. Extension rods for use in spraying trees and vines may be obtained for either. For operations on a very small scale a good hand-syringe may be used, but as a general thing it will be best to invest a few dollars more and get a small tank sprayer, as this throws a continuous stream or spray and holds a much larger amount of the spraying solution. Whatever type is procured, get a brass machine it will out-wear three or four of those made of cheaper metal, which succumbs very quickly to the, corroding action of the strong poisons and chemicals used in them.

Of implements for harvesting, beside the spade, prong-hoe and spading- fork, very few are used in the small garden, as most of them need not only long rows to be economically used, but horse- power also. The onion harvester attachment

for the double wheel hoe may be used with advantage in loosening onions, beets, turnips, etc., from the soil or for cutting spinach. Running the hand-plow close on either side of carrots, parsnips and other deep-growing vegetables will aid materially in getting them out. For fruit picking, with tall trees, the wire-fingered fruit-picker, secured to the end of a long handle, will be of great assistance, but with the modern method of using low-headed trees it will not be needed.

Another class of garden implements are those used in pruning but where this is attended to properly from the start, a good sharp jack-knife and a pair of pruning shears will easily handle all the work of the kind necessary.

Still another sort of garden device is that used for supporting the plants; such as stakes, trellises, wires, etc. Altogether too little attention usually is given these, as with proper care in storing over winter they will not only last for years, but add greatly to the convenience of cultivation and to the neat appearance of the garden.

As a final word to the intending purchaser of garden tools, I would say: first thoroughly investigate the different sorts available, and when buying, do not forget that a good tool or a well-made machine will be giving you satisfactory use long, long after the price is forgotten, while a poor one is a constant source of discomfort. Get good tools, and take good care of them. And let me repeat that a few dollars a year, judiciously spent, for tools afterward well cared for, will soon give you a very complete set, and add to your garden profit and pleasure.

GARDEN PESTS.

If we could garden without any interference from the pests which attack plants, then indeed gardening would be a simple matter. But all the time we must watch out for these little foes little in size, but tremendous in the havoc they make.

As human illness may often be prevented by healthful conditions, so pests may be kept away by strict garden cleanliness. Heaps of waste is lodging places for the breeding of insects. I do not think a compost pile will do the harm, but unkempt, uncared-for spots seem to invite trouble.

There are certain helps to keeping pests down. The constant stirring up of the soil by earthworms is an aid in keeping the soil open to air and water. Many of our common birds feed upon insects. The sparrows, robins, chickadees, meadow larks and orioles are all examples of birds that help in this way. Some insects feed on other and harmful insects. Some kinds of ladybugs do this good deed. The ichneumon-fly helps too. And toads are wonders in the number of insects they can consume at one meal. The toad deserves very kind treatment from all of us.

Each gardener should try to make her or his garden into a place attractive to birds and toads. A good birdhouse, grain sprinkled about in early spring, a water-place, are invitations for birds to stay a while in your garden. If you wish toads, fix things up for them too. During a hot summer day a toad likes to rest in the shade. By night he is ready to go forth to eat but not to kill, since toads prefer live food. How can one "fix up" for toads? Well, one thing to do is to prepare a retreat, quiet, dark and damp. A few stones of some size underneath the shade of a shrub with perhaps a carpeting of damp leaves would appear very fine to a toad.

There are two general classes of insects known by the way they do their work. One kind gnaws at the plant really taking pieces of it into its system. This kind of

insect has a mouth fitted to do this work. Grasshoppers and caterpillars are of this sort. The other kind sucks the juices from a plant. This, in some ways, is the worst sort. Plant lice belong here, as do mosquitoes, which prey on us. All the scale insects fasten themselves on plants, and suck out the life of the plants.

Now can we fight these chaps? The gnawing fellows may be caught with poison sprayed upon plants, which they take into their bodies with the plant. The Bordeaux mixture which is a poison sprayed upon plants for this purpose.

In the other case the only thing is to attack the insect direct. So certain insecticides, as they are called, are sprayed on the plant to fall upon the insect. They do a deadly work of attacking, in one way or another, the body of the insect.

Sometimes we are much troubled with underground insects at work. You have seen a garden covered with ant hills. Here is a remedy, but one of which you must be careful.

This question is constantly being asked, 'How can I tell what insect is doing the destructive work?' Well, you can tell partly by the work done, and partly by seeing the insect itself. This latter thing is not always so easy to accomplish. I had cutworms one season and never saw one. I saw only the work done. If stalks of tender plants are cut clean off be pretty sure the cutworm is abroad. What does he look like? Well, that is a hard question because his family is a large one. Should you see sometime a grayish striped caterpillar, you may know it is a cutworm. But because of its habit of resting in the ground during the day and working by night, it is difficult to catch sight of one. The cutworm is around early in the season ready to cut the flower stalks of the hyacinths. When the peas come on a bit later, he is ready for them. A very good way to block him off is to put paper collars, or tin ones, about the plants. These collars should be about an inch away from the plant.

Of course, plant lice are more common. Those we see are often green in color. But they may be red, yellow or brown. Lice are easy enough to find since they are always clinging to their host. As sucking insects they have to cling close to a plant for food, and one is pretty sure to find them. But the biting insects do their work, and then go hide. That makes them much more difficult to deal with.

Rose slugs do great damage to the rose bushes. They eat out the body of the leaves, so that just the veining is left. They are soft-bodied, green above and yellow below.

A beetle, the striped beetle, attacks young melons and squash leaves. It eats the leaf by riddling out holes in it. This beetle, as its name implies, is striped. The back is black with yellow stripes running lengthwise.

Then there are the slugs, which are garden pests. The slug will devour almost any garden plant, whether it is a flower or a vegetable. They lay lots of eggs in old rubbish heaps. Do you see the good of cleaning up rubbish? The slugs do more harm in the garden than almost any other single insect pest. You can discover them in the following way. There is a trick for bringing them to the surface of the ground in the day time. You see they rest during the day below ground. So just water the soil in which the slugs are supposed to be. How are you to know where they are? They are quite likely to hide near the plants they are feeding on. So water the ground with some nice clean lime water. This will disturb them, and up they'll poke to see what the matter is.

Beside these most common of pests, pests which attack many kinds of plants, there are special pests for special plants. Discouraging, is it not? Beans have pests of their own; so have potatoes and cabbages. In fact, the vegetable garden has many inhabitants. In the flower garden lice are very bothersome, the cutworm and the slug have a good time there, too, and ants often get very numerous as the season advances. But for real discouraging insect troubles the

vegetable garden takes the prize. If we were going into fruit to any extent, perhaps the vegetable garden would have to resign in favor of the fruit garden.

A common pest in the vegetable garden is the tomato worm. This is a large yellowish or greenish striped worm. Its work is to eat into the young fruit.

A great, light green caterpillar is found on celery. This caterpillar may be told by the black bands, one on each ring or segment of its body.

The squash bug may be told by its brown body, which is long and slender, and by the disagreeable odor from it when killed. The potato bug is another fellow to look out for. It is a beetle with yellow and black stripes down its crusty back. The little green cabbage worm is a perfect nuisance. It is a small caterpillar and smaller than the tomato worm. These are perhaps the most common of garden pests by name.

LANDSCAPE GARDENING.

Landscape gardening has often been likened to the painting of a picture. Your art-work teacher has doubtless told you that a good picture should have a point of chief interest, and the rest of the points simply go to make more beautiful the central idea, or to form a fine setting for it. So in landscape gardening there must be in the gardener's mind a picture of what he desires the whole to be when he completes his work.

From this study we shall be able to work out a little theory of landscape gardening.

Let us go to the lawn. A good extent of open lawn space is always beautiful. It is restful. It adds a feeling of space to even small grounds. So we might generalize and say that it is well to keep open lawn spaces. If one covers his lawn space with many trees, with little flower beds here and there, the general effect is choppy and fussy. It is a bit like an over-dressed person. One's grounds lose all individuality thus treated. A single tree or a small group is not a bad arrangement on the lawn. Do not centre the tree or trees. Let them drop a bit into the background. Make a pleasing side feature of them. In choosing trees one must keep in mind a number of things. You should not choose an overpowering tree; the tree should be one of good shape, with something interesting about its bark, leaves, flowers or fruit. While the poplar is a rapid grower, it sheds its leaves early and so is left standing, bare and ugly, before the fall is old. Mind you, there are places where a row or double row of Lombardy poplars is very effective. But I think you'll agree with me that one lone poplar is not. The catalpa is quite lovely by itself. Its leaves are broad, its flowers attractive, the seed pods which cling to the tree until away into the winter, add a bit of picture siveness. The bright berries of the ash, the brilliant foliage of the sugar maple, the blossoms of the tulip tree, the bark of the white birch, and the leaves of the copper beech all these are beauty points to consider.

Place makes a difference in the selection of a tree. Suppose the lower portion of the grounds is a bit low and moist, then the spot is ideal for a willow. Don't group trees together which look awkward. A long-looking poplar does not go with a nice rather rounded little tulip tree. A juniper, so neat and prim, would look silly beside a spreading chestnut. One must keep proportion and suitability in mind.

I'd never advise the planting of a group of evergreens close to a house, and in the front yard. The effect is very gloomy indeed. Houses thus surrounded are over capped by such trees and are not only gloomy to live in, but truly unhealthful. The chief requisite inside a house is sunlight and plenty of it.

As trees are chosen because of certain good points, so shrubs should be. In a clump I should wish some which bloomed early, some which bloomed late, and some for the beauty of their fall foliage, some for the color of their bark and others for the fruit. Some spireas and the forsythia bloom early. The red bark of the dogwood makes a bit of color all winter, and the red berries of the barberry cling to the shrub well into the winter.

Certain shrubs are good to use for hedge purposes. A hedge is rather prettier usually than a fence. The Californian privet is excellent for this purpose. Osage orange, Japan barberry, buckthorn, Japan quince, and Van Houtte's spirea are other shrubs which make good hedges.

I forgot to say that in tree and shrub selection it is usually better to choose those of the locality one lives in. Unusual and foreign plants do less well and often harmonize but poorly with their new setting.

Landscape gardening may follow along very formal lines or along informal lines. The first would have straight paths, straight rows in stiff beds, everything, as the name tells, perfectly formal. The other method is, of course, the exact opposite. There are danger points in each.

The formal arrangement is likely to look too stiff; the informal, too fussy, too wiggly. As far as paths go, keep this in mind, that a path should always lead somewhere. That is its business to direct one to a definite place. Now, straight, even paths are not displeasing if the effect is to be that of a formal garden. The danger in the curved path is an abrupt curve, a whirligig effect. It is far better for you to stick to straight paths unless you can make a really beautiful curve. No one can tell you how to do this.

Garden paths may be of gravel, of dirt, or of grass. One sees grass paths in some very lovely gardens. I doubt, however, if they would serve as well in your

small gardens. Your garden areas are so limited that they should be re-spaded each season, and the grass paths are a great bother in this work. Of course, a gravel path makes a fine appearance, but again you may not have gravel at your command. It is possible for any of you to dig out the path for two feet. Then put in six inches of stone or clinker. Over this, pack in the dirt, rounding it slightly toward the centre of the path. There should never be depressions through the central part of paths, since these form convenient places for water to stand. The under layer of stone makes a natural drainage system.

A building often needs the help of vines or flowers or both to tie it to the grounds in such a way as to form a harmonious whole. Vines lend themselves well to this work. It is better to plant a perennial vine, and so let it form a permanent part of your landscape scheme. The Virginia creeper, wistaria, honeysuckle, a climbing rose, the clematis and trumpet vine are all most satisfactory.

Close your eyes and picture a house of natural color, that mellow gray of the weathered shingles. Now add to this old house a purple wistaria. Can you see the beauty of it? I shall not forget soon a rather ugly corner of my childhood home, where the dining room and kitchen met. Just there climbing over, and falling over a trellis was a trumpet vine. It made beautiful an awkward angle, an ugly bit of carpenter work.

Of course, the morning-glory is an annual vine, as is the moon-vine and wild cucumber. Now, these have their special function. For often, it is necessary to cover an ugly thing for just a time, until the better things and better times come. The annual is 'the chap' for this work.

Along an old fence a hop vine is a thing of beauty. One might try to rival the woods' landscape work. For often one sees festooned from one rotted tree to another the ampelopsis vine.

Flowers may well go along the side of the building, or bordering a walk. In general, though, keep the front lawn space open and unbroken by beds. What lovelier in early spring than a bed of daffodils close to the house? Hyacinths and tulips, too, form a blaze of glory. These are little or no bother, and start the spring aright. One may make of some bulbs an exception to the rule of unbroken front lawn. Snowdrops and crocuses planted through the lawn are beautiful. They do not disturb the general effect, but just blend with the whole. One expert bulb gardener says to take a basketful of bulbs in the fall, walk about your grounds, and just drop bulbs out here and there. Wherever the bulbs drop, plant them. Such small bulbs as those we plant in lawns should be in groups of four to six. Daffodils may be thus planted, too. You all remember the grape hyacinths that grow all through Katharine's side yard.

The place for a flower garden is generally at the side or rear of the house. The backyard garden is a lovely idea, is it not? Who wishes to leave a beautiful looking front yard, turn the corner of a house, and find a dump heap? Not I. The flower garden may be laid out formally in neat little beds, or it may be more of a careless, hit-or-miss sort. Both have their good points. Great masses of bloom are attractive.

You should have in mind some notion of the blending of color. Nature appears not to consider this at all, and still gets wondrous effects. This is because of the tremendous amount of her perfect background of green, and the limitlessness of her space, while we are confined at the best to relatively small areas. So we should endeavor not to blind people's eyes with clashes of colors which do not at close range blend well. In order to break up extremes of colors you can always use masses of white flowers, or something like mignonette, which is in effect green.

Finally, let us sum up our landscape lesson. The grounds are a setting for the house or buildings. Open, free lawn spaces, a tree or a proper group well placed

flowers which do not clutter up the front yard, groups of shrubbery these are points to be remembered. The paths should lead somewhere, and be either straight or well curved. If one starts with a formal garden, one should not mix the informal with it before the work is done.

THE CULTIVATION OF VEGETABLES.

Before taking up the garden vegetables individually, I shall outline the general practice of cultivation, which applies to all.

The purposes of cultivation are three to get rid of weeds, and to stimulate growth by (1) letting air into the soil and freeing unavailable plant food, and (2) by conserving moisture.

As to weeds, the gardener of any experience need not be told the importance of keeping his crops clean. He has learned from bitter and costly experience the price of letting them get anything resembling a start. He knows that one or two days' growth, after they are well up, followed perhaps by a day or so of rain, may easily double or treble the work of cleaning a patch of onions or carrots, and that where weeds have attained any size they cannot be taken out of sowed crops without doing a great deal of injury. He also realizes, or should, that every day's growth means just so much available plant food stolen from under the very roots of his legitimate crops.

Instead of letting the weeds get away with any plant food, he should be furnishing more, for clean and frequent cultivation will not only break the soil up mechanically, but let in air, moisture and heat all essential in effecting those chemical changes necessary to convert non- available into available plant food. Long before the science in the case was discovered, the soil cultivators had

learned by observation the necessity of keeping the soil nicely loosened about their growing crops. Even the lanky and untutored aborigine saw to it that his squaw not only put a bad fish under the hill of maize but plied her shell hoe over it. Plants need to breathe. Their roots need air. You might as well expect to find the rosy glow of happiness on the wan cheeks of a cotton-mill child slave as to expect to see the luxuriant dark green of healthy plant life in a suffocated garden.

Important as the question of air is, that of water ranks beside it. You may not see at first what the matter of frequent cultivation has to do with water. But let us stop a moment and look into it. Take a strip of blotting paper, dip one end in water, and watch the moisture run up hill, and soak up through the blotter. The scientists have labeled that "capillary attraction" the water crawls up little invisible tubes formed by the texture of the blotter. Now take a similar piece, cut it across, hold the two cut edges firmly together, and try it again. The moisture refuses to cross the line: the connection has been severed.

In the same way the water stored in the soil after a rain begins at once to escape again into the atmosphere. That on the surface evaporates first, and that which has soaked in begins to soak in through the soil to the surface. It is leaving your garden, through the millions of soil tubes, just as surely as if you had a two-inch pipe and a gasoline engine, pumping it into the gutter night and day! Save your garden by stopping the waste. It is the easiest thing in the world to do cut the pipe in two. By frequent cultivation of the surface soil not more than one or two inches deep for most small vegetables the soil tubes are kept broken, and a mulch of dust is maintained. Try to get over every part of your garden, especially where it is not shaded, once in every ten days or two weeks. Does that seem like too much work? You can push your wheel hoe through, and thus keep the dust mulch as a constant protection, as fast as you can walk. If you wait for the weeds, you will nearly have to crawl through, doing more or less harm by disturbing your growing plants, losing all the plant food (and they will take the cream) which they have consumed, and actually putting in more hours of infinitely

more disagreeable work. If the beginner at gardening has not been convinced by the facts given, there is only one thing left to convince him experience.

Having given so much space to the reason for constant care in this matter, the question of methods naturally follows. Get a wheel hoe. The simplest sorts will not only save you an infinite amount of time and work, but do the work better, very much better than it can be done by hand. You can grow good vegetables, especially if your garden is a very small one, without one of these labor-savers, but I can assure you that you will never regret the small investment necessary to procure it.

With a wheel hoe, the work of preserving the soil mulch becomes very simple. If one has not a wheel hoe, for small areas very rapid work can be done with the scuffle hoe.

The matter of keeping weeds cleaned out of the rows and between the plants in the rows is not so quickly accomplished. Where hand-work is necessary, let it be done at once. Here are a few practical suggestions that will reduce this work to a minimum, (1) Get at this work while the ground is soft; as soon as the soil begins to dry out after a rain is the best time. Under such conditions the weeds will pull out by the roots, without breaking off. (2) Immediately before weeding, go over the rows with a wheel hoe, cutting shallow, but just as close as possible, leaving a narrow, plainly visible strip which must be hand-weeded. The best tool for this purpose is the double wheel hoe with disc attachment, or hoes for large plants. (3) See to it that not only the weeds are pulled but that every inch of soil surface is broken up. It is fully as important that the weeds just sprouting be destroyed, as that the larger ones be pulled up. One stroke of the weeder or the fingers will destroy a hundred weed seedlings in less time than one weed can be pulled out after it gets a good start. (4) Use one of the small hand-weeders until you become skilled with it. Not only may more work be done but the fingers will be saved unnecessary wear.

The skilful use of the wheel hoe can be acquired through practice only. The first thing to learn is that it is necessary to watch the wheels only: the blades, disc or rakes will take care of themselves.

The operation of "hilling" consists in drawing up the soil about the stems of growing plants, usually at the time of second or third hoeing. It used to be the practice to hill everything that could be hilled "up to the eyebrows," but it has gradually been discarded for what is termed "level culture"; and you will readily see the reason, from what has been said about the escape of moisture from the surface of the soil; for of course the two upper sides of the hill, which may be represented by an equilateral triangle with one side horizontal, give more exposed surface than the level surface represented by the base. In wet soils or seasons hilling may be advisable, but very seldom otherwise. It has the additional disadvantage of making it difficult to maintain the soil mulch which is so desirable.

Rotation of crops.

There is another thing to be considered in making each vegetable do its best, and that is crop rotation, or the following of any vegetable with a different sort at the next planting.

With some vegetables, such as cabbage, this is almost imperative, and practically all are helped by it. Even onions, which are popularly supposed to be the proving exception to the rule, are healthier, and do as well after some other crop, provided the soil is as finely pulverized and rich as a previous crop of onions would leave it.

Here are the fundamental rules of crop rotation:

(1) Crops of the same vegetable, or vegetables of the same family (such as turnips and cabbage) should not follow each other.

(2) Vegetables that feed near the surface, like corn, should follow deep-rooting crops.

(3) Vines or leaf crops should follow root crops.

(4) Quick-growing crops should follow those occupying the land all season.

These are the principles which should determine the rotations to be followed in individual cases. The proper way to attend to this matter is when making the planting plan. You will then have time to do it properly, and will need to give it no further thought for a year.

With the above suggestions in mind, and put to use, it will not be difficult to give the crops those special attentions which are needed to make them do their very best.

VEGETABLE CULTURE.

As a rule, we choose to grow bush beans rather than pole beans. I cannot make up my mind whether or not this is from sheer laziness. In a city backyard the tall varieties might perhaps be a problem since it would be difficult to get poles. But these running beans can be trained along old fences and with little urging will run up the stalks of the tallest sunflowers. So that settles the pole question. There is an ornamental side to the bean question. Suppose you plant these tall beans at the extreme rear end of each vegetable row. Make arches with supple tree limbs, binding them over to form the arch. Train the beans over these. When one stands facing the garden, what a beautiful terminus these bean arches make.

Beans like rich, warm, sandy soil. In order to assist the soil is sure to dig deeply, and work it over thoroughly for bean culture. It never does to plant beans before the world has warmed up from its spring chills. There is another advantage in early digging of soil. It brings to the surface eggs and larvae of insects. The birds eager for food will even follow the plough to pick from the soil these choice morsels. A little lime worked in with the soil is helpful in the cultivation of beans.

Bush beans are planted in drills about eighteen inches apart, while the pole-bean rows should be three feet apart. The drills for the bush limas should be further apart than those for the other dwarf beans say three feet. This amount of space gives opportunity for cultivation with the hoe. If the running beans climb too high just pinch off the growing extreme end and this will hold back the upward growth.

Among bush beans are the dwarf, snap or string beans, the wax beans, the bush limas, one variety of which is known as brittle beans. Among the pole beans are the pole limas, wax and scarlet runner. The scarlet runner is a beauty for decorative effects. The flowers are scarlet and are fine against an old fence. These are quite lovely in the flower garden. Where one wishes a vine, this is good to plant for one gets both a vegetable, bright flowers and a screen from the

one plant. When planting beans put the bean in the soil edgewise with the eye down.

Beets like rich, sandy loam, also. Fresh manure worked into the soil is fatal for beets, as it is for many other crops. But we will suppose that nothing is available but fresh manure. Some gardeners say to work this into the soil with great care and thoroughness. But even so, there is danger of a particle of it getting next to a tender beet root. The following can be done; Dig a trench about a foot deep, spread a thin layer of manure in this, cover it with soil, and plant above this. By the time the main root strikes down to the manure layer, there will be little harm done. Beets should not be transplanted. If the rows are one foot apart there is ample space for cultivation. Whenever the weather is really settled, then these seeds may be planted. Young beet tops make fine greens. Greater care should be taken in handling beets than usually is shown. When beets are to be boiled, if the tip of the root and the tops are cut off, the beet bleeds. This means a loss of good material. Pinching off such parts with the fingers and doing this not too closely to the beet itself is the proper method of handling.

There are big coarse members of the beet and cabbage families called the mangel wurzel and ruta бага. About here these are raised to feed to the cattle. They are a great addition to a cow's dinner.

The cabbage family is a large one. There is the cabbage proper, then cauliflower, broccoli or a more hardy cauliflower, kale, Brussels sprouts and kohlrabi, a cabbage-turnip combination.

Cauliflower is a kind of refined, high-toned cabbage relative. It needs a little richer soil than cabbage and cannot stand the frost. A frequent watering with manure water gives it the extra richness and water it really needs. The outer leaves must be bent over, as in the case of the young cabbage, in order to get the white head. The dwarf varieties are rather the best to plant.

Kale is not quite so particular a cousin. It can stand frost. Rich soil is necessary and early spring planting, because of slow maturing. It may be planted in September for early spring work.

Brussels sprouts are a very popular member of this family. On account of their size many people who do not like to serve poor, common old cabbage will serve these. Brussels sprouts are interesting in their growth. The plant stalk runs skyward. At the top, umbrella like, is a close head of leaves, but this is not what we eat. Shaded by the umbrella and packed all along the stalk are delicious little cabbages or sprouts. Like the rest of the family a rich soil is needed and plenty of water during the growing period. The seed should be planted in May, and the little plants transplanted into rich soil in late July. The rows should be eighteen inches apart, and the plants one foot apart in the rows.

Kohlrabi is a go-between in the families of cabbage and turnip. It is sometimes called the turnip-root cabbage. Just above the ground the stem of this plant swells into a turnip-like vegetable. In the true turnip the swelling is underground, but like the cabbage, kohlrabi forms its edible part above ground. It is easy to grow. Only it should develop rapidly, otherwise the swelling gets woody, and so loses its good quality. Sow out as early as possible; or sow inside in March and transplant to the open. Plant in drills about two feet apart. Set the plants about one foot apart or thin out to this distance. To plant one hundred feet of drill buy half an ounce of seed. Seed goes a long way, you see. Kohlrabi is served and prepared like turnip. It is a very satisfactory early crop.

Before leaving the cabbage family I should like to say that the cabbage called Savoy is an excellent variety to try. It should always have an early planting under cover, say in February, and then be transplanted into open beds in March or April. If the land is poor where you are to grow cabbage, then by all means choose Savoy.

Carrots are of two general kinds: those with long roots, and those with short roots. If long-rooted varieties are chosen, then the soil must be worked down to a depth of eighteen inches, surely. The shorter ones will do well in eight inches of well-worked sandy soil. Do not put carrot seed into freshly manured land. Another point in carrot culture is one concerning the thinning process. As the little seedlings come up you will doubtless find that they are much, much too close together. Wait a bit, thin a little at a time, so that young, tiny carrots may be used on the home table. These are the points to jot down about the culture of carrots.

The cucumber is the next vegetable in the line. This is a plant from foreign lands. Some think that the cucumber is really a native of India. A light, sandy and rich soil is needed I mean rich in the sense of richness in organic matter. When cucumbers are grown outdoors, as we are likely to grow them, they are planted in hills. Nowadays, they are grown in hothouses; they hang from the roof, and are a wonderful sight. In the greenhouse a hive of bees is kept so that cross-fertilization may go on.

But if you intend to raise cucumbers follow these directions: Sow the seed inside, cover with one inch of rich soil. In a little space of six inches diameter, plant six seeds. Place like a bean seed with the germinating end in the soil. When all danger of frost is over, each set of six little plants, soil and all, should be planted in the open. Later, when danger of insect pests is over, thin out to three plants in a hill. The hills should be about four feet apart on all sides.

Before the time of Christ, lettuce was grown and served. There is a wild lettuce from which the cultivated probably came. There are a number of cultivated vegetables which have wild ancestors, carrots, turnips and lettuce being the most common among them. Lettuce may be tucked into the garden almost anywhere. It is surely one of the most decorative of vegetables. The compact head, the

green of the leaves, the beauty of symmetry all these are charming characteristics of lettuces.

As the summer advances and as the early sowings of lettuce get old they tend to go to seed. Don't let them. Pull them up. None of us are likely to go into the seed-producing side of lettuce. What we are interested in is the raising of tender lettuce all the season. To have such lettuce in mid and late summer is possible only by frequent plantings of seed. If seed is planted every ten days or two weeks all summer, you can have tender lettuce all the season. When lettuce gets old it becomes bitter and tough.

Melons are most interesting to experiment with. We suppose that melons originally came from Asia, and parts of Africa. Melons are a summer fruit. Over in England we find the muskmelons often grown under glass in hothouses. The vines are trained upward rather than allowed to lie prone. As the melons grow large in the hot, dry atmosphere, just the sort which is right for their growth, they become too heavy for the vine to hold up. So they are held by little bags of netting, just like a tennis net in size of mesh. The bags are supported on nails or pegs. It is a very pretty sight I can assure you. Over here usually we raise our melons outdoors. They are planted in hills. Eight seeds are placed two inches apart and an inch deep. The hills should have a four foot sweep on all sides; the watermelon hills ought to have an allowance of eight to ten feet. Make the soil for these hills very rich. As the little plants get sizeable say about four inches in height reduce the number of plants to two in a hill. Always in such work choose the very sturdiest plants to keep. Cut the others down close to or a little below the surface of the ground. Pulling up plants is a shocking way to get rid of them. I say shocking because the pull is likely to disturb the roots of the two remaining plants. When the melon plant has reached a length of a foot, pinch off the end of it. This pinch means this to the plant: just stop growing long, take time now to grow branches. Sand or lime sprinkled about the hills tends to keep bugs away.

The word pumpkin stands for good, old-fashioned pies, for Thanksgiving, for grandmother's house. It really brings more to mind than the word squash. I suppose the squash is a bit more useful, when we think of the fine Hubbard, and the nice little crooked-necked summer squashes; but after all, I like to have more pumpkins. And as for Jack-o'-lanterns why they positively demand pumpkins. In planting these, the same general directions hold good which were given for melons. And use these same for squash-planting, too. But do not plant the two cousins together, for they have a tendency to run together. Plant the pumpkins in between the hills of corn and let the squashes go in some other part of the garden.

WILD-FLOWER GARDEN.

A wild-flower garden has a most attractive sound. One thinks of long tramps in the woods, collecting material, and then of the fun in fixing up a real for sure wild garden.

Many people say they have no luck at all with such a garden. It is not a question of luck, but a question of understanding, for wild flowers are like people and each has its personality. What a plant has been accustomed to in Nature it desires always. In fact, when removed from its own sort of living conditions, it sickens and dies. That is enough to tell us that we should copy Nature herself. Suppose you are hunting wild flowers. As you choose certain flowers from the woods, notice the soil they are in, the place, conditions, the surroundings, and the neighbors.

Suppose you find dog-tooth violets and wind-flowers growing near together. Then place them so in your own new garden. Suppose you find a certain violet enjoying an open situation; then it should always have the same. You see the

point, do you not? If you wish wild flowers to grow in a tame garden make them feel at home. Cheat them into almost believing that they are still in their native haunts.

Wild flowers ought to be transplanted after blossoming time is over. Take a trowel and a basket into the woods with you. As you take up a few, a columbine, or a hepatica, is sure to take with the roots some of the plant's own soil, which must be packed about it when replanted.

The bed into which these plants are to go should be prepared carefully before this trip of yours. Surely you do not wish to bring those plants back to wait over a day or night before planting. They should go into new quarters at once. The bed needs soil from the woods, deep and rich and full of leaf mold. The under drainage system should be excellent. Then plants are not to go into water-logged ground. Some people think that all wood plants should have a soil saturated with water. But the woods themselves are not water-logged. It may be that you will need to dig your garden up very deeply and put some stone in the bottom. Over this the top soil should go. And on top, where the top soil once was, put a new layer of the rich soil you brought from the woods.

Before planting water the soil well. Then as you make places for the plants put into each hole some of the soil which belongs to the plant which is to be put there.

I think it would be a rather nice plan to have a wild-flower garden giving a succession of bloom from early spring to late fall; so let us start off with March, the hepatica, spring beauty and saxifrage. Then comes April bearing in its arms the beautiful columbine, the tiny bluets and wild geranium. For May there are the dog-tooth violet and the wood anemone, false Solomon's seal, Jack-in-the-pulpit, wake robin, bloodroot and violets. June will give the bellflower, mullein, bee balm and foxglove. I would choose the gay butterfly weed for July. Let turtle head,

aster, Joe Pye weed, and Queen Anne's lace make the rest of the season brilliant until frost.

Let us have a bit about the likes and dislikes of these plants. After you are once started you'll keep on adding to this wild-flower list.

There is no one who doesn't love the hepatica. Before the spring has really decided to come, this little flower pokes its head up and puts all else to shame. Tucked under a covering of dry leaves the blossoms wait for a ray of warm sunshine to bring them out. These embryo flowers are further protected by a fuzzy covering. This reminds one of a similar protective covering which new fern leaves have. In the spring a hepatica plant wastes no time on getting a new suit of leaves. It makes its old ones do until the blossom has had its day. Then the new leaves, started to be sure before this, have a chance. These delayed, are ready to help out next season. You will find hepaticas growing in clusters, sort of family groups. They are likely to be found in rather open places in the woods. The soil is found to be rich and loose. So these should go only in partly shaded places and under good soil conditions. If planted with other woods specimens give them the benefit of a rather exposed position, that they may catch the early spring sunshine. I should cover hepaticas over with a light litter of leaves in the fall. During the last days of February, unless the weather is extreme take this leaf covering away. You'll find the hepatica blossoms all ready to poke up their heads.

The spring beauty hardly allows the hepatica to get ahead of her. With a white flower which has dainty tracings of pink, a thin, wiry stem, and narrow, grass-like leaves, this spring flower cannot be mistaken. You will find spring beauties growing in great patches in rather open places. Plant a number of the roots and allow the sun good opportunity to get at them. For this plant loves the sun.

The other March flower mentioned is the saxifrage. This belongs in quite a different sort of environment. It is a plant which grows in dry and rocky places.

Often one will find it in chinks of rock. There is an old tale to the effect that the saxifrage roots twine about rocks and work their way into them so that the rock itself splits. Anyway, it is a rock garden plant. I have found it in dry, sandy places right on the borders of a big rock. It has white flower clusters borne on hairy stems.

The columbine is another plant that is quite likely to be found in rocky places. Standing below a ledge and looking up, one sees nestled here and there in rocky crevices one plant or more of columbine. The nodding red heads bob on wiry, slender stems. The roots do not strike deeply into the soil; in fact, often the soil hardly covers them. Now, just because the columbine has little soil, it does not signify that it is indifferent to the soil conditions. For it always has lived, and always should live, under good drainage conditions. I wonder if it has struck you, how really hygienic plants are? Plenty of fresh air, proper drainage, and good food are fundamentals with plants.

It is evident from study of these plants how easy it is to find out what plants like. After studying their feelings, then do not make the mistake of huddling them all together under poor drainage conditions.

I always have a feeling of personal affection for the bluets. When they come I always feel that now things are beginning to settle down outdoors. They start with rich, lovely, little delicate blue blossoms. As June gets hotter and hotter their color fades a bit, until at times they look quite worn and white. Some people call them Quaker ladies, others innocence. Under any name they are charming. They grow in colonies, sometimes in sunny fields, sometimes by the road-side. From this we learn that they are more particular about the open sunlight than about the soil.

If you desire a flower to pick and use for bouquets, then the wild geranium is not your flower. It droops very quickly after picking and almost immediately drops its

petals. But the purplish flowers are showy, and the leaves, while rather coarse, are deeply cut. This latter effect gives certain boldness to the plant that is rather attractive. The plant is found in rather moist, partly shaded portions of the woods. I like this plant in the garden. It adds good color and permanent color as long as blooming time lasts, since there is no object in picking it.

There are numbers and numbers of wild flowers I might have suggested. These I have mentioned were not given for the purpose of a flower guide, but with just one end in view your understanding of how to study soil conditions for the work of starting a wild-flower garden.

If you fear results, take but one or two flowers and study just what you select. Having mastered, or better, become acquainted with a few, add more another year to your garden. I think you will love your wild garden best of all before you are through with it. It is a real study, you see.

Recommended Resources

1) [**Bonsai Gardening Secrets.**](#)

Discover over 95 pages of insider secrets to creating stunning bonsai trees.

2) [**Home And Garden - Country And Rural Life.**](#)

Gardening and Birds, Raising Chickens and Goats, Baking Bread.

3) [**The Weekend Gardener.**](#)

The Busy Persons Guide to a Beautiful Backyard Garden!

4) [**The Home Of Indoor Plant Care Info.**](#)

We provide indoor plant care information for the homeowner, garden center personnel, and plantscape technicians.

5) [**Growing & Enjoying Your Own Rose Garden.**](#)

Everything you need to know to get your own rose garden established and thriving.

6) **[Square Watermelon Secrets Revealed!](#)**

Instant buzz home garden eBook. Author grew his own square watermelon and shows how to. Its great fun and curiosity.

7) **[Organic Profits.](#)**

How to make money with Organic Gardening.

8) **[Your Perfect Lawn.](#)**

Step by step guide on building and maintaining a perfect garden lawn.

9) **[Garden's Moles.](#)**

Tricks and secrets to rid of them.

10) **[The Comfortable Lazy Garden.](#)**

Gardening and landscaping with native plants.

How To Get Traffic, Sales and Profits You DESERVE...

For the first time ever, Raam Anand reveals the 15 'secret' strategies that he used to get over 65117 REAL visitors to his website in 30 days!

Get a first-hand look at how you will be able to become a "TRAFFIC MAGNET" and attract unlimited, highly targeted TRAFFIC to YOUR website, including the amazing traffic "Blueprint" that you can use TODAY to flood your website with visitors.



Visit <http://Traffic-Sales-Profits.com/> to learn how you can **magically increase YOUR traffic, sales and profits** like Raam.

